

The Psychosocial Impact of Lung Cancer-Related Stigma: Findings from the Cancer Experience Registry

Joanne S Buzaglo PhD¹, Melissa F Miller PhD MPH¹, Alexandra K Zaleta PhD¹, James Johnson MPH CPH¹, Linda House RN BSN MSM¹, Emily K Porensky PhD¹, Janine Cataldo RN PhD FAAN²

¹Cancer Support Community, Research and Training Institute, Philadelphia PA; ²University of California San Francisco, San Francisco, CA

Background

- Patients with lung cancer (LC) report greater psychosocial distress and depression than people with other types of cancers.
- Patients with LC also report experiencing stigmatization because the disease is associated with smoking.

Aims

- The objectives of this study were to 1) characterize patients' experience of lung cancer stigma, and 2) examine the relationships between distress, depression, and lung cancer stigma.

Methods

- 47 LC patients enrolled in the online Cancer Experience Registry answered questions about their cancer history, 13 LC stigma-related statements (0 = *strongly disagree* to 4 = *strongly agree*), and 27 cancer-related distress items (0 = *not at all concerned* to 4 = *very seriously concerned*) based on CancerSupportSource®. Distress questions included a 4-item depression risk subscale.
- Analytic procedures included: descriptive statistics, independent samples t-tests and bivariate correlations between stigma-related statements and clinical and demographic variables (age, gender, and LC type), multivariate linear regression with stigma statements predicting distress (square root of sum of 27 items), and logistic regression with stigma statements predicting risk for depression (score ≥ 6). Regressions were adjusted for potentially confounding variables (LC type, age, gender).

Participants

	N = 47	Median / n	SD / %
Age		64 Range: 39 - 79	
Non-Hispanic White		40	85%
Time since diagnosis (years)		2 Range: <1 - 13	
Lung cancer type			
Non-small cell (NSCLC)		38	84%
Small cell (SCLC)		7	16%
Experienced recurrence		16	34%
History of metastatic disease		25	53%

Acknowledgments

This work is sponsored by Lilly Oncology, Boehringer Ingelheim, Bristol-Meyers Squibb, and AbbVie.

References

- Zabora et al. (2001). The prevalence of psychological distress by cancer site. *Psycho-Oncology*, 10: 19-28.
- Chambers et al. (2012). A systematic review of stigma and nihilism on lung cancer outcomes. *BMC Cancer*, 12, 184.
- Miller et al. (2014). Discriminatory power of a 25-item distress screening tool: A cross-sectional survey of 251 cancer survivors. *Qual Life Res*, 23(10): 2855-63.

Results

Proportion of Patients Agreeing with Lung Cancer Stigma Statements



- Most participants agreed that some people assume LC is caused by smoking even if the person with cancer never smoked (91%), and that lung cancer is viewed as a self-inflicted disease (74%).
- Of note, 81% reported being open to speaking with people about their lung cancer diagnosis, and 83% believed their health care team treated them as well as people with other cancer diagnoses.

Lung Cancer Stigma Experience Differs by Cancer Type, Gender, and Age

	Mean		t	p
Lung Cancer Type	NSCLC	SCLC		
LC is viewed as self-inflicted disease	3.1	1.9	2.60	.013*
Open to speaking about LC	3.4	2.6	2.11	.041*
People uncomfortable b/c associate LC with death	2.5	1.6	2.04	.048*
Feel guilty because of having LC	1.5	2.7	-2.15	.037*
Try to keep LC a secret	0.7	1.7	-2.54	.015*

Gender

	Mean	t	p	
LC is viewed as self-inflicted disease	2.3	3.2	-2.46	.018*
Worry about people discriminating against me	0.8	1.7	-2.43	.019*
Open to speaking about LC	2.8	3.4	-2.09	.042*

Age

	r	p
Worry about people discriminating against me	-.36	.026*
LC is viewed as self-inflicted disease	-.33	.045*

Note: *p < .05

- Patients with NSCLC** were more likely to believe that lung cancer is viewed as a self-inflicted disease and that some people are uncomfortable around them because they associate lung cancer with death. They were also more open to speaking with people about their lung cancer diagnosis.
- Patients with SCLC** felt more guilt because of having lung cancer and tried harder to keep their diagnosis a secret.
- Younger patients and females** worried more about people discriminating against them and agreed more strongly that lung cancer is viewed as a self-inflicted disease. Females were also more open than males to speaking with people about their lung cancer diagnosis.

Lung Cancer Stigma is Associated with Elevated Distress

Stigma-related statement	β (SE)*	p
I feel guilty because of having LC	0.46 (0.19)	.004
I am open to speaking about LC	-0.37 (0.32)	.031
I try to keep LC a secret	0.49 (0.29)	.004
I worry people may judge me	0.43 (0.21)	.007
I stopped socializing because of reactions	0.40 (0.21)	.012

Note: The square root of the overall summary score for distress was used to normalize the distribution.
*Adjusted for LC type, age, and gender

- Patients with greater distress tended to feel more guilt because of having lung cancer, tried harder to keep lung cancer a secret, worried more that people may judge them, and reported socializing less due to reactions people had to their lung cancer diagnosis.
- Patients with lower levels of distress were more open to speaking with people about their lung cancer diagnosis.

Lung Cancer Stigma is Associated with Risk for Depression

Stigma-related statement	Adjusted odds ratio (95% CI)*	p
I feel guilty because of having LC	2.28 (1.12, 4.67)	.024

Note: Risk for depression defined as score ≥6 on the depression subscale (37% of the sample).
*Adjusted for LC type, age, and gender

- Patients who reported more guilt about having lung cancer were at greater risk for clinical levels of depression.

Implications and Conclusions

- As advances in lung cancer treatment emerge and people living with LC live longer, the short- and long-term implications of living with stigma become increasingly important.
- This study adds to the evidence that stigma is associated with greater distress and greater risk for clinically-significant depression in lung cancer patients.
- Our findings suggest that the type of lung cancer (NSCLC vs. SCLC) is related to the kinds of feelings patients experience related to stigma. Additionally, younger age and female gender are specifically associated with greater endorsement of stigma.
- These results support the development and evaluation of interventions to reduce the impact of lung-cancer related stigma, especially guilt, and the risks it confers for cancer-related distress and depression.